

GESTATIONAL DIABETES
 PRETERM DELIVERY
 POSTPARTUM SCREENING
 LONG TERM SURVEILLANCE
 RISK STRATIFICATION
 SMALL-FOR-GESTATIONAL-AGE BIRTH
 PRECLAMPSIA
 GESTATIONAL HYPERTENSION


10 KICKS
 OBSTETRICS
www.10KICKS.com

Gestational Care

10 KICKS
 OBSTETRICS

ask your obstetrician about

- Baby Kick Count
- Hypertension
- Diabetes
- Hypoxia



Mother & Child
 Safety



Baby Kick Count

10KICKS

O B S T E T R I C S

Online Connected Maternity Stillbirth Risk remote monitor baby safety

Incidence

In 2015 there were 2.6 million stillbirths globally, with more than 7178 deaths a day. The of these deaths occurred in developing countries. Ninety-eight percent majority occurred in low- and middle-income countries. About half of all stillbirths occur in the intrapartum period, representing the greatest time of risk. Estimated proportion of stillbirths that are intrapartum varies from 10% in developed regions to 59% in south Asia.

https://www.who.int/maternal_child_adolescent/epidemiology/stillbirth/en/

Gestational Hypertension
Gestational Diabetes
Gestational Hypoxia
Gestational BMI

Diabetes

Children who are exposed to high blood glucose in the womb are at higher risk of developing Type II Diabetes later in life”


<https://www.diabetes.ie/living-with-diabetes/diabetes-pregnancy/gestational-diabetes/>



Health & Medical Devices separate order; manual or wireless

Mother & Child Safety

© 10KICKS. Copyright. All Rights Reserved



Abdominal Circumference
Head Circumference
Femur Length
Ovulation Calculator

Maternal Mental Health Pre-Post Childbirth DEPRESSION



DOWNLOAD APP



10KICKS
O B S T E T R I C S
Division Of IMEUS
www.10KICKS.com